

Model of hematopoiesis with structured cell differentiation

Vlastimil Severa

Masaryk university in Brno

Podlesí, 11.-13.9.2012



MINISTERSTVO ŠKOLSTVÍ,
MLÁDEŽE A TĚLOVÝCHOVY



INVESTICE DO ROZVOJE VZDĚLÁVÁNÍ



- 1 Introduction
- 2 Algorithmization of hematopoiesis
- 3 The model of common strange cells pool
- 4 Simulations

Process of hematopoiesis

- Hematopoiesis is a process taking place in bone marrow.

Process of hematopoiesis

- Hematopoiesis is a process taking place in bone marrow.
- Hematopoietic stem cells (HSC) give rise to all blood cells through differentiation and cell division.

Process of hematopoiesis

- Hematopoiesis is a process taking place in bone marrow.
- Hematopoietic stem cells (HSC) give rise to all blood cells through differentiation and cell division.
- The count of new cells is controled by negative feedback of actual size of blood cells populations.


Process of hematopoiesis

- Hematopoiesis is a process taking place in bone marrow.
- Hematopoietic stem cells (HSC) give rise to all blood cells through differentiation and cell division.
- The count of new cells is controled by negative feedback of actual size of blood cells populations.
- Due to the processes of control delays occur.

Process of hematopoiesis

- Hematopoiesis is a process taking place in bone marrow.
- Hematopoietic stem cells (HSC) give rise to all blood cells through differentiation and cell division.
- The count of new cells is controled by negative feedback of actual size of blood cells populations.
- Due to the processes of control delays occur.
- Hematopoietic diseases: Chronic myelogenous leukemia, Cyclical neutropenia, etc.

Scheme of hematopoiesis



Cell characteristics

We observe two characteristics of blood cells: *level and commitment*.

Level indicates *stemness* (or on the other hand *maturation*) of the cell.

Commitment is qualitative characteristic indicating which lineage the cell has presumption to differentiate in.

Cell characteristics

We observe two characteristics of blood cells: *level and commitment*.

Level indicates *stemness* (or on the other hand *maturation*) of the cell.

Commitment is qualitative characteristic indicating which lineage the cell has presumption to differentiate in.

- 0 ... any commitment
- L , M ... commitment to lymphoid, or myeloid lineage
- M_E , M_{MEG} , M_{GM} ... myeloid cells commitment to erythroid, thrombocytic, or leukocytic lineage
- M_G , M_M ... leukocyt commitment to granulocytic, or monocytic lineage

Cell characteristics

We observe two characteristics of blood cells: *level and commitment*.

Level indicates *stemness* (or on the other hand *maturation*) of the cell.

Commitment is qualitative characteristic indicating which lineage the cell has presumption to differentiate in.

- 0 ... any commitment
- L , M ... commitment to lymphoid, or myeloid lineage
- M_E , M_{MEG} , M_{GM} ... myeloid cells commitment to erythroid, thrombocytic, or leukocytic lineage
- M_G , M_M ... leukocyte commitment to granulocytic, or monocytic lineage

Then $\mathcal{X}_{i,c}$ stands for the cell of level i and commitment c .

Strange cells pool

Strange cells pool is a group of cells in which commitment arise or can be changed. Assume there are three strange cells pools:

Strange cells pool


Strange cells pool is a group of cells in which commitment arise or can be changed. Assume there are three strange cells pools:

- Cells without commitment specify to the cells with commitment L and M . There are cells on level 1 to $k - 1$.
- Cells with commitment M specify to the cells with commitment M_E , M_{MEG} and M_{GM} . There are cells on level k to $l - 1$.
- Cells with commitment M_{GM} specify to the cells with commitment M_G and M_M . There are cells on level l to $m - 1$.

Strange cells pool

Strange cells pool is a group of cells in which commitment arise or can be changed. Assume there are three strange cells pools:

- Cells without commitment specify to the cells with commitment L and M . There are cells on level 1 to $k - 1$.
- Cells with commitment M specify to the cells with commitment M_E , M_{MEG} and M_{GM} . There are cells on level k to $l - 1$.
- Cells with commitment M_{GM} specify to the cells with commitment M_G and M_M . There are cells on level l to $m - 1$.



Cell processes

Cell division

$$\mathcal{X}_{i,c} \begin{cases} \nearrow \mathcal{X}_{i,c_1} \\ \searrow \mathcal{X}_{i,c_1} \end{cases}$$

$$\mathcal{X}_{i,c} \begin{cases} \nearrow \mathcal{X}_{i+1,c_1} \\ \searrow \mathcal{X}_{i+1,c_1} \end{cases}$$

$$\mathcal{X}_{i,c} \begin{cases} \nearrow \mathcal{X}_{i,c_1} \\ \searrow \mathcal{X}_{i,c_2} \end{cases}$$

$$\mathcal{X}_{i,c} \begin{cases} \nearrow \mathcal{X}_{i+1,c_1} \\ \searrow \mathcal{X}_{i+1,c_2} \end{cases}$$

$$\mathcal{X}_{i,c} \begin{cases} \nearrow \mathcal{X}_{i,c_1} \\ \searrow \mathcal{X}_{i+1,c_1} \end{cases}$$

$$\mathcal{X}_{i,c} \begin{cases} \nearrow \mathcal{X}_{i,c_1} \\ \searrow \mathcal{X}_{i+1,c_2} \end{cases}$$

Cell processes

Cell division

$$\mathcal{X}_{i,c} \begin{cases} \nearrow \mathcal{X}_{i,c_1} \\ \searrow \mathcal{X}_{i,c_1} \end{cases}$$

$$\mathcal{X}_{i,c} \begin{cases} \nearrow \mathcal{X}_{i,c_1} \\ \searrow \mathcal{X}_{i,c_2} \end{cases}$$

$$\mathcal{X}_{i,c} \begin{cases} \nearrow \mathcal{X}_{i,c_1} \\ \searrow \mathcal{X}_{i+1,c_1} \end{cases}$$

$$\mathcal{X}_{i,c} \begin{cases} \nearrow \mathcal{X}_{i+1,c_1} \\ \searrow \mathcal{X}_{i+1,c_1} \end{cases}$$

$$\mathcal{X}_{i,c} \begin{cases} \nearrow \mathcal{X}_{i+1,c_1} \\ \searrow \mathcal{X}_{i+1,c_2} \end{cases}$$

$$\mathcal{X}_{i,c} \begin{cases} \nearrow \mathcal{X}_{i,c_1} \\ \searrow \mathcal{X}_{i+1,c_2} \end{cases}$$

Cell transformation

$$\mathcal{X}_{i,c_1} \longrightarrow \mathcal{X}_{i,c_2}$$

$$\mathcal{X}_{i,c_1} \longrightarrow \mathcal{X}_{i+1,c_1}$$

$$\mathcal{X}_{i,c_1} \longrightarrow \mathcal{X}_{i+1,c_2}$$

Cell processes

Cell division

$$\mathcal{X}_{i,c} \begin{cases} \nearrow \mathcal{X}_{i,c_1} \\ \searrow \mathcal{X}_{i,c_1} \end{cases}$$

$$\mathcal{X}_{i,c} \begin{cases} \nearrow \mathcal{X}_{i,c_1} \\ \searrow \mathcal{X}_{i,c_2} \end{cases}$$

$$\mathcal{X}_{i,c} \begin{cases} \nearrow \mathcal{X}_{i,c_1} \\ \searrow \mathcal{X}_{i+1,c_1} \end{cases}$$

$$\mathcal{X}_{i,c} \begin{cases} \nearrow \mathcal{X}_{i+1,c_1} \\ \searrow \mathcal{X}_{i+1,c_1} \end{cases}$$

$$\mathcal{X}_{i,c} \begin{cases} \nearrow \mathcal{X}_{i+1,c_1} \\ \searrow \mathcal{X}_{i+1,c_2} \end{cases}$$

$$\mathcal{X}_{i,c} \begin{cases} \nearrow \mathcal{X}_{i,c_1} \\ \searrow \mathcal{X}_{i+1,c_2} \end{cases}$$

$$Q_{(j_1, c_1), (j_2, c_2)}^{(i, c)}$$

stands for the probability that the cell $\mathcal{X}_{i,c}$ divides to cells \mathcal{X}_{j_1, c_1} and \mathcal{X}_{j_2, c_2}

Cell transformation

$$\mathcal{X}_{i,c_1} \longrightarrow \mathcal{X}_{i,c_2}$$

$$\mathcal{X}_{i,c_1} \longrightarrow \mathcal{X}_{i+1,c_1}$$

$$\mathcal{X}_{i,c_1} \longrightarrow \mathcal{X}_{i+1,c_2}$$

Cell processes

Cell division

$$\mathcal{X}_{i,c} \begin{cases} \nearrow \mathcal{X}_{i,c_1} \\ \searrow \mathcal{X}_{i,c_1} \end{cases}$$

$$\mathcal{X}_{i,c} \begin{cases} \nearrow \mathcal{X}_{i+1,c_1} \\ \searrow \mathcal{X}_{i+1,c_1} \end{cases}$$

$$\mathcal{X}_{i,c} \begin{cases} \nearrow \mathcal{X}_{i,c_1} \\ \searrow \mathcal{X}_{i,c_2} \end{cases}$$

$$\mathcal{X}_{i,c} \begin{cases} \nearrow \mathcal{X}_{i+1,c_1} \\ \searrow \mathcal{X}_{i+1,c_2} \end{cases}$$

$$\mathcal{X}_{i,c} \begin{cases} \nearrow \mathcal{X}_{i,c_1} \\ \searrow \mathcal{X}_{i+1,c_1} \end{cases}$$

$$\mathcal{X}_{i,c} \begin{cases} \nearrow \mathcal{X}_{i,c_1} \\ \searrow \mathcal{X}_{i+1,c_2} \end{cases}$$

$$Q_{(j_1, c_1), (j_2, c_2)}^{(i, c)}$$

stands for the probability that the cell $\mathcal{X}_{i,c}$ divides to cells \mathcal{X}_{j_1, c_1} and \mathcal{X}_{j_2, c_2}

Cell transformation

$$\mathcal{X}_{i, c_1} \longrightarrow \mathcal{X}_{i, c_2}$$

$$\mathcal{X}_{i, c_1} \longrightarrow \mathcal{X}_{i+1, c_1}$$

$$\mathcal{X}_{i, c_1} \longrightarrow \mathcal{X}_{i+1, c_2}$$

$$P_{(j, c_2)}^{(i, c_1)}$$

stands for the probability that the cell \mathcal{X}_{i, c_1} transforms to cell \mathcal{X}_{j, c_2}

Processes assumptions

Assume that:

- Probabilities $P_{(j,c_2)}^{(i,c_1)}$ and $Q_{(j_1,c_1),(j_2,c_2)}^{(i,c)}$ which aren't showed are equal to zero.

Processes assumptions

Assume that:

- Probabilities $P_{(j,c_2)}^{(i,c_1)}$ and $Q_{(j_1,c_1),(j_2,c_2)}^{(i,c)}$ which aren't showed are equal to zero.
- Apoptosis (*cell death*) is possible, but not sure. That for the cell $\mathcal{X}_{(i,c)}$ means

$$0 < \sum_{\iota} \sum_{\kappa \in C} P_{(\iota,\kappa)}^{(i,c)} + \sum_{\iota,\lambda} \sum_{\kappa,\delta \in C} Q_{(\iota,\kappa),(\lambda,\delta)}^{(i,c)} < 1,$$

where $C = \{0, L, M, M_E, M_{MEG}, M_{GM}, M_G, M_M\}$.

Processes assumptions

Assume that:

- Probabilities $P_{(j,c_2)}^{(i,c_1)}$ and $Q_{(j_1,c_1),(j_2,c_2)}^{(i,c)}$ which aren't showed are equal to zero.
- Apoptosis (*cell death*) is possible, but not sure. That for the cell $\mathcal{X}_{(i,c)}$ means

$$0 < \sum_{\iota} \sum_{\kappa \in C} P_{(\iota,\kappa)}^{(i,c)} + \sum_{\iota,\lambda} \sum_{\kappa,\delta \in C} Q_{(\iota,\kappa),(\lambda,\delta)}^{(i,c)} < 1,$$

where $C = \{0, L, M, M_E, M_{MEG}, M_{GM}, M_G, M_M\}$.

- With growing level the probability of commitment change decreases, or

$$P_{(j,c_2)}^{(i,c_1)} \geq P_{(j+1,c_2)}^{(i+1,c_1)},$$


$$Q_{(i+1,c_1),(i+1,c_2)}^{(i+1,c_1)} \leq Q_{(i,c_2),(i,c_2)}^{(i,c_1)} \leq Q_{(i,c_1),(i,c_2)}^{(i,c_1)} \leq Q_{(i,c_1),(i,c_1)}^{(i,c_1)}.$$

Model scheme

Let $x_{i,c}$ stands for the group of all cells $\mathcal{X}_{i,c}$.

Model scheme

Let $x_{i,c}$ stands for the group of all cells $\mathcal{X}_{i,c}$.



Model equations

$$x'_{0,0}(t) = -G_{0,0}x_{0,0}(t) + V_{(0,0)}^{(0,0)}x_{0,0}(t - \tau_{0,0}),$$

$$x'_{1,0}(t) = -G_{1,0}x_{1,0}(t) + V_{(1,0)}^{(1,0)}x_{1,0}(t - \tau_{1,0}) + V_{(1,0)}^{(0,0)}x_{0,0}(t - \tau_{0,0}),$$

$$x'_{2,c_1}(t) = -G_{2,c_1}x_{2,c_1}(t) + V_{(2,c_1)}^{(2,c_1)}x_{2,c_1}(t - \tau_{2,c_1}) + V_{(2,c_1)}^{(2,c_2)}x_{2,c_2}(t - \tau_{2,c_2}) \\ + V_{(2,c_1)}^{(1,0)}x_{1,0}(t - \tau_{1,0}),$$

$$x'_{i,c_1}(t) = -G_{i,c_1}x_{i,c_1}(t) + V_{(i,c_1)}^{(i,c_1)}x_{i,c_1}(t - \tau_{i,c_1}) + V_{(i,c_1)}^{(i,c_2)}x_{i,c_2}(t - \tau_{i,c_2}) \\ + V_{(i,c_1)}^{(i-1,c_1)}x_{i-1,c_1}(t - \tau_{i-1,c_1}) + V_{(i,c_1)}^{(i-1,c_2)}x_{i-1,c_2}(t - \tau_{i-1,c_2}),$$

$$x'_{k-1,c_1}(t) = -G_{k-1,c_1}x_{k-1,c_1}(t) + V_{(k-1,c_1)}^{(k-1,c_1)}x_{k-1,c_1}(t - \tau_{k-1,c_1}) \\ + V_{(k-1,c_1)}^{(k-1,c_2)}x_{k-1,c_2}(t - \tau_{k-1,c_2}) + V_{(k-1,c_1)}^{(k-2,c_1)}x_{k-2,c_1}(t - \tau_{k-2,c_1}) \\ + V_{(k-1,c_1)}^{(k-2,c_2)}x_{k-2,c_2}(t - \tau_{k-2,c_2}) - \Lambda_L x_{k-1,c_1}(t),$$

where $i = 3, \dots, k-2$ and $c_1, c_2 \in \{L, M\}$, $c_1 \neq c_2$.

Coefficients $G_{i,c}$

Assume cell processes coefficients as functions in the form

$$P_{(j,c_t)}^{(i,c)} = \mu_{(j,c_t)}^{(i,c)} \frac{\left(\theta_{(j,c_t)}^{(i,c)}\right)^n}{\left(\theta_{(j,c_t)}^{(i,c)}\right)^n + x_{j,c_t}^n}, \quad Q_{(j_1,c_1)(j_2,c_2)}^{(i,c)} = \mu_{(j_1,c_1)(j_2,c_2)}^{(i,c)} \frac{\left(\theta_{(j_1,c_1)(j_2,c_2)}^{(i,c)}\right)^n}{\left(\theta_{(j_1,c_1)(j_2,c_2)}^{(i,c)}\right)^n + (x_{j_1,c_1} x_{j_2,c_2})^n}.$$

Coefficients $G_{i,c}$

Assume cell processes coefficients as functions in the form

$$P_{(j,c_t)}^{(i,c)} = \mu_{(j,c_t)}^{(i,c)} \frac{\left(\theta_{(j,c_t)}^{(i,c)}\right)^n}{\left(\theta_{(j,c_t)}^{(i,c)}\right)^n + x_{j,c_t}^n}, \quad Q_{(j_1,c_1)(j_2,c_2)}^{(i,c)} = \mu_{(j_1,c_1)(j_2,c_2)}^{(i,c)} \frac{\left(\theta_{(j_1,c_1)(j_2,c_2)}^{(i,c)}\right)^n}{\left(\theta_{(j_1,c_1)(j_2,c_2)}^{(i,c)}\right)^n + (x_{j_1,c_1} x_{j_2,c_2})^n}.$$

Next we consider function

$$H_{i,c} = \sum_{\iota} \sum_{\kappa \in C} P_{(\iota,\kappa)}^{(i,c)} + \sum_{\iota,\lambda} \sum_{\kappa,\delta \in C} Q_{(\iota,\kappa),(\lambda,\delta)}^{(i,c)}.$$

Coefficients $G_{i,c}$

Assume cell processes coefficients as functions in the form

$$P_{(j,c_t)}^{(i,c)} = \mu_{(j,c_t)}^{(i,c)} \frac{\left(\theta_{(j,c_t)}^{(i,c)}\right)^n}{\left(\theta_{(j,c_t)}^{(i,c)}\right)^n + x_{j,c_t}^n}, \quad Q_{(j_1,c_1)(j_2,c_2)}^{(i,c)} = \mu_{(j_1,c_1)(j_2,c_2)}^{(i,c)} \frac{\left(\theta_{(j_1,c_1)(j_2,c_2)}^{(i,c)}\right)^n}{\left(\theta_{(j_1,c_1)(j_2,c_2)}^{(i,c)}\right)^n + (x_{j_1,c_1} x_{j_2,c_2})^n}.$$

Next we consider function

$$H_{i,c} = \sum_{\iota} \sum_{\kappa \in C} P_{(\iota,\kappa)}^{(i,c)} + \sum_{\iota,\lambda} \sum_{\kappa,\delta \in C} Q_{(\iota,\kappa),(\lambda,\delta)}^{(i,c)}.$$

And finally we can define coefficients (functions) used in the system of equations

$$G_{i,c} = M_{i,c} \frac{H_{i,c}^n}{\Theta_{i,c}^n + H_{i,c}^n}.$$

Coefficients V_{j,c_2}^{i,c_1}

During the processes cells can be lost with rate γ_{i,c_1} , i.e. at the end there is overall

$$G_{i,c_1} x_{i,c_1} e^{-\gamma_{i,c_1} \tau_{i,c_1}}.$$

Coefficients V_{j,c_2}^{i,c_1}

During the processes cells can be lost with rate γ_{i,c_1} , i.e. at the end there is overall

$$G_{i,c_1} x_{i,c_1} e^{-\gamma_{i,c_1} \tau_{i,c_1}}.$$

Cells that have completed the processes which last for time τ_{i,c_1} are distributed into cells groups. It is appropriate that the cells distribution corresponds to rates between the cell processes coefficients.

Coefficients V_{j,c_2}^{i,c_1}

During the processes cells can be lost with rate γ_{i,c_1} , i.e. at the end there is overall

$$G_{i,c_1} x_{i,c_1} e^{-\gamma_{i,c_1} \tau_{i,c_1}}.$$

Cells that have completed the processes which last for time τ_{i,c_1} are distributed into cells groups. It is appropriate that the cells distribution corresponds to rates between the cell processes coefficients.

So we set

$$V_{j,c_2}^{i,c_1} = \left(2Q_{(j,c_2),(j,c_2)}^{(i,c_1)} + \sum_{\iota} \sum_{\kappa \in C} Q_{(j,c_2),(\iota,\kappa)}^{(i,c_1)} + P_{(j,c_2)}^{(i,c_1)} \right) \frac{G_{i,c_1}}{H_{i,c_1}} e^{-\gamma_{i,c_1} \tau_{i,c_1}},$$

where $j \neq \iota$ or $c_2 \neq \kappa$.

Origin of parameters

Parameter values were chosen according to some papers dealing with modelling of hematopoiesis, mainly by Michael C. Mackey [1, 2, 3].

Origin of parameters

Parameter values were chosen according to some papers dealing with modelling of hematopoiesis, mainly by Michael C. Mackey [1, 2, 3].

For parameters $\gamma_{i,c}$ it seems the most important one is $\gamma_{0,0}$. Even the small change disturbs stability of the model. For $i > 1$ $\gamma_{i,c}$ has to change significantly and there only occurs decrease in cells quantity.

Origin of parameters

Parameter values were chosen according to some papers dealing with modelling of hematopoiesis, mainly by Michael C. Mackey [1, 2, 3].

For parameters $\gamma_{i,c}$ it seems the most important one is $\gamma_{0,0}$. Even the small change disturbs stability of the model. For $i > 1$ $\gamma_{i,c}$ has to change significantly and there only occurs decrease in cells quantity.

For delay $\tau_{i,c}$ we obtain quite similar results.

Origin of parameters


Parameter values were chosen according to some papers dealing with modelling of hematopoiesis, mainly by Michael C. Mackey [1, 2, 3].

For parameters $\gamma_{i,c}$ it seems the most important one is $\gamma_{0,0}$. Even the small change disturbs stability of the model. For $i > 1$ $\gamma_{i,c}$ has to change significantly and there only occurs decrease in cells quantity.

For delay $\tau_{i,c}$ we obtain quite similar results.


Generally we can say that most important are changes for cell groups $x_{0,0}$ and $x_{1,0}$. For higher levels there is small effect on the whole population. But this presumption has to be investigated on more complex model.

Parameters $\gamma_{i,c}$




$\gamma_{0,0} = 0.0526$, $\gamma_{1,0} = 0.186$, $\gamma_{i,c} = 0.0027$ for $i > 1$

Parameters $\gamma_{i,c}$




$\gamma_{0,0} = 0.0526$, $\gamma_{1,0} = 0.186$, $\gamma_{i,c} = 0.0027$ for $i > 1$




$\gamma_{0,0} = 0.0526$, $\gamma_{i,c} = 0.0027$ for $i > 0$


Parameters $\gamma_{i,c}$



$\gamma_{0,0} = 0.0526$, $\gamma_{1,0} = 0.186$, $\gamma_{i,c} = 0.0027$ for $i > 1$




$\gamma_{0,0} = 0.0526$, $\gamma_{i,c} = 0.0027$ for $i > 0$




$\gamma_{i,c} = 0.0027$ for all i


Parameters $\gamma_{i,c}$




$\gamma_{0,0} = 0.0526$, $\gamma_{1,0} = 0.186$, $\gamma_{i,c} = 0.0027$ for $i > 1$



$\gamma_{0,0} = 0.0526$, $\gamma_{i,c} = 0.0027$ for $i > 0$




$\gamma_{i,c} = 0.0027$ for all i




$\gamma_{i,c} = 0.0027$ for all i

Parameters $\Theta_{i,c}$




$\Theta_{i,c} = 0.2$ for all i

Parameters $\Theta_{i,c}$




$\Theta_{i,c} = 0.2$ for all i




$\Theta_{0,0} = 0.6, \Theta_{i,c} = 0.2$ for $i > 0$


Parameters $\Theta_{i,c}$



$\Theta_{i,c} = 0.2$ for all i




$\Theta_{0,0} = 0.6, \Theta_{i,c} = 0.2$ for $i > 0$




$\Theta_{1,0} = 0.6, \Theta_{i,c} = 0.2$ for $i = 1$ or $i > 0$

Parameters $\Theta_{i,c}$




$\Theta_{0,0} = 0.4, \Theta_{1,0} = 0.6, \Theta_{i,c} = 0.2$ for $i > 1$

Parameters $\Theta_{i,c}$




$\Theta_{0,0} = 0.4, \Theta_{1,0} = 0.6, \Theta_{i,c} = 0.2$ for $i > 1$




$\Theta_{0,0} = 0.4, \Theta_{1,0} = 0.6, \Theta_{i,c} = 0.2$ for $i > 1$

Parameters $\Theta_{i,c}$




$\Theta_{0,0} = 0.4$, $\Theta_{1,0} = 0.6$, $\Theta_{i,c} = 0.2$ for $i > 1$



$\Theta_{0,0} = 0.4$, $\Theta_{1,0} = 0.6$, $\Theta_{i,c} = 0.2$ for $i > 1$

Compare that with simulation in [3].
There is shown model for platelet
population in collie with cyclical
neutropenia under treatment.



Bibliography



Michael C. Mackey: Unified Hypothesis for the Origin of Aplastic Anemia and Periodic Hematopoiesis, *Blood*, **Vol.51**, pp. 941–956, 1978



Caroline Colijn, Michael C. Mackey: A mathematical model of hematopoiesis: I. Periodic chronic myelogenous leukemia, *Journal of Theoretical Biology*, **Vol. 237**, pp. 117–132, 2005



Caroline Colijn, Michael C. Mackey: A mathematical model of hematopoiesis: II. Cyclical neutropenia, *Journal of Theoretical Biology*, **Vol. 237**, pp. 133–146, 2005



Mostafa Adimy, Fabien Crauste, Abderrahim El Abdallaoui: Asymptotic Behavior of a Discrete Maturity Structured System of Hematopoietic Stem Cell Dynamics with Several Delays, *Mathematical Modelling of Natural Phenomena*, **Vol.2**, pp. 1–19, 2007.

Thank you for your attention.