

SEMI-ABELIAN MONADIC CATEGORIES

MARINO GRAN AND JIŘÍ ROSICKÝ *

ABSTRACT. We characterize semi-abelian monadic categories and their localizations. These results are then used to obtain a characterization of pointed protomodular quasimonadic categories, and in particular of protomodular quasivarieties.

INTRODUCTION

The notion of semi-abelian category can be considered as intermediate between the notion of Barr-exact category and the one of abelian category. Semi-abelian categories were introduced by Janelidze, Marki and Tholen [12] in a closed connection with the more general protomodular categories due to Bourn [4]. A finitely complete category with a zero object is protomodular if and only if it satisfies the split short five lemma. Semi-abelian categories are defined as exact protomodular categories with finite coproducts and a zero object. These categories are suitable to develop several basic aspects of homological algebra of groups and rings [6], as well as an abstract theory of commutators and of ideals [5]. Among the examples of semi-abelian categories there are the categories of groups, rings, commutative rings, Lie algebras, Boolean algebras, crossed modules and compact Hausdorff groups.

Every variety of universal algebras is an exact category, and abelian varieties are precisely those whose theories contain abelian group operations 0 , $-$ and $+$ in such a way that these operations are homomorphisms. When the theory of a variety \mathcal{V} only contains group operations 0 , $-$ and $+$, then \mathcal{V} is semi-abelian. Bourn and Janelidze recently characterized semi-abelian varieties [7] as those whose theories contain a unique constant 0 , binary operations $\alpha_0, \alpha_1, \dots, \alpha_{n-1}$ for $n \geq 1$ and a $(n+1)$ -ary operation β satisfying the equations $\alpha_i(x, x) = 0$ for $i = 0, 1, \dots, n-1$ and $\beta(\alpha_0(x, y), \alpha_1(x, y), \dots, \alpha_{n-1}(x, y), y) = x$. The case $n = 1$ shows that the above-mentioned existence of a group operation suffices to guarantee that \mathcal{V} is semi-abelian, by setting $\alpha_0(x, y) = x - y$ and $\beta(x, y) = x + y$. Varieties of algebras satisfying these axioms have been also studied by Ursini [19], who called them classically ideal determined.

In the present paper we show that the characterization of semi-abelian varieties can be extended to infinitary and many sorted ones. As a consequence we prove that C^* -algebras form a semi-abelian category, and we provide explicit operations witnessing this fact. Every variety of infinitary many-sorted algebras is exact and locally presentable. Having a general exact locally presentable category \mathcal{C} , we can consider a varietal hull \mathcal{V} of \mathcal{C} with respect to a chosen regular generator G of \mathcal{C} .

Key words and phrases. Semi-abelian categories, monadic categories, varieties, quasivarieties, localizations.

2000 Mathematics Subject Classification. 18C20, 18C35, 08C05, 08C15, 18E10, 18E35.

* Work supported by the Grant Agency of the Czech Republic under the grant 201/02/0148 and by the Université du Littoral Côte d'Opale.

Following [21] the category \mathcal{C} is a localization of \mathcal{V} and, thus, \mathcal{C} is semi-abelian whenever \mathcal{V} is semi-abelian. Using our characterization of semi-abelian infinitary many-sorted varieties we get a characterization of their localizations. In particular, any exact locally presentable category \mathcal{C} containing a regular generator G which is a cogroup in \mathcal{C} is semi-abelian.

In the last section we characterize protomodular quasimonadic categories, and then protomodular quasivarieties and their localizations.

1. SEMI-ABELIAN MONADIC CATEGORIES

Let us recall that a functor $U: \mathcal{V} \rightarrow \mathit{Sets}$ is monadic if it has a left adjoint F and the comparison functor from \mathcal{V} to the category of algebras $\mathit{Alg}(T)$ of the induced monad $T = UF$ is an equivalence. A category \mathcal{V} is monadic over sets if there exists a monadic functor $U: \mathcal{V} \rightarrow \mathit{Sets}$. We shall often use the term “monadic category” to indicate a monadic category over sets. Monadic categories are precisely those given by a class of single-sorted infinitary operations and a class of equations such that free algebras exist [14]. Free algebras always exist if \mathcal{V} is determined by a set of operations and a set of equations. In any case, the elements of $UF(n)$ (where n is a cardinal and $F(n)$ is a free algebra over n) correspond to n -ary terms.

The first result we are going to prove is a straightforward generalization of the characterization of semi-abelian varieties given in [7]. We shall follow the presentation given in [3].

Let us recall that in any finitely complete pointed category the *split short five lemma* means the following statement: given a diagram (1)

$$\begin{array}{ccccc}
 A' & \xrightarrow{\quad k' \quad} & B' & \xleftarrow{\quad s' \quad} & C' \\
 \downarrow f & & \downarrow g & \xrightarrow{\quad p' \quad} & \downarrow h \\
 A & \xrightarrow{\quad k \quad} & B & \xleftarrow{\quad s \quad} & C \\
 & & & \xrightarrow{\quad p \quad} &
 \end{array}$$

where all squares are commutative, $p \circ s = 1_C$, $p' \circ s' = 1_{C'}$, $k = \ker(p)$, $k' = \ker(p')$ and f and h are isomorphisms, then g is an isomorphism.

Since a monadic category over sets is always exact and cocomplete, it is *semi-abelian* exactly when it is pointed and it satisfies the split short five lemma.

1.1. Theorem. *Let $U: \mathcal{V} \rightarrow \mathit{Set}$ be a monadic functor. Then \mathcal{V} is semi-abelian if and only if the corresponding theory has a unique constant 0, binary terms α_i $i \in n$, where $n \geq 1$ is a cardinal, and a $(n+1)$ -ary term β satisfying the equations*

$$\alpha_i(x, x) = 0 \quad \text{for } i \in n$$

and

$$\beta(\alpha_0(x, y), \alpha_1(x, y), \dots, \alpha_i(x, y), \dots, y) = x$$

Proof. Let $F(x, y)$ and $F(y)$ be the free algebras on $\{x, y\}$ and $\{y\}$ respectively, and let $p: F(x, y) \rightarrow F(y)$ be the homomorphism determined by $p(x) = p(y) = y$. Then p is split by the inclusion $s: F(y) \rightarrow F(x, y)$. Let $k: K \rightarrow F(x, y)$ be a kernel of p and A a subalgebra of $F(x, y)$ generated by $UK \cup UF(y)$. Since the codomain restriction of k is a kernel of the domain restriction $p': A \rightarrow F(y)$ of p , $A = F(x, y)$

by the split short five lemma. Hence there are elements $k_i \in K$, with $i \in n$, (where $n \geq 1$ is a cardinal) and a $(n+1)$ -ary term β such that $x = \beta(k_0, k_1, \dots, k_i, \dots, y)$. Since $k_i \in K$, there are binary terms $\alpha_i(x, y)$ such that $\alpha_i(x, y) = k_i$ for $i \in n$. Moreover, one obviously has that $\alpha_i(x, x) = 0$.

Conversely, assume that the terms satisfying the conditions in the theorem exist. Let us then consider the diagram (1), and we are going to prove that the arrow g is an isomorphism.

First consider a and b in B' such that $g(a) = g(b)$. Then

$$(h \circ p')(\alpha_i(a, b)) = (p \circ g)(\alpha_i(a, b)) = p(\alpha_i(g(a), g(b))) = p(0) = 0$$

and then $\alpha_i(a, b)$ is in A' for $i \in n$. Since $(k \circ f)(\alpha_i(a, b)) = 0$, it follows that $\alpha_i(a, b) = 0$ for $i \in n$, which implies that $a = b$ because

$$b = \beta(\alpha_0(b, b), \alpha_1(b, b), \dots, b) = \beta(0, 0, \dots, b) = \beta(\alpha_0(a, b), \alpha_1(a, b), \dots, b) = a.$$

Consequently, g is injective.

In order to check that g is surjective, let us consider any $b \in B$. We define $a = (s' \circ h^{-1} \circ p)(b)$. We have

$$p(\alpha_i(b, g(a))) = \alpha_i(p(b), (p \circ g)(a)) = \alpha_i(p(b), p(b)) = 0.$$

Hence $\alpha_i(b, g(a))$ is in A for $i \in n$ and then

$$b = \beta(\alpha_0(b, g(a)), \alpha_1(b, g(a)), \dots, g(a)) = \beta(g(a_0), g(a_1), \dots, g(a)) = g(\beta(a_0, a_1, \dots, a))$$

where $f(a_i) = \alpha_i(b, g(a))$ for $i \in n$. Thus g is surjective. \square

1.2. Remark. a) The same argument applies to varieties of S -sorted algebras, i.e. to monadic categories $U: \mathcal{V} \rightarrow \text{Set}^S$. One just needs terms α_i for $i \in n$ and β in each sort $s \in S$.

b) A similar argument allows one to characterize protomodular monadic categories: one simply has to replace the single constant 0 by constants e_i for $i \in n$, with the properties $\alpha_i(x, x) = e_i$ and one keeps the axiom

$$\beta(\alpha_0(x, y), \alpha_1(x, y), \dots, \alpha_i(x, y), \dots, y) = x.$$

The following well-known simple lemma [6] immediately follows from the definitions:

1.3. Lemma. *Let $H: \mathcal{C} \rightarrow \mathcal{L}$ be a conservative pullback preserving functor, where \mathcal{C} and \mathcal{L} are pointed categories with pullbacks and \mathcal{L} satisfies the split short five lemma. Then \mathcal{C} satisfies the split short five lemma.*

1.4. Example. Let \mathcal{C} be the category of non-unital C^* -algebras, where arrows are continuous homomorphisms of involutive algebras. The forgetful functor from \mathcal{C} to the category of involutive algebras preserves finite limits and reflects isomorphisms (see 1.3.3 and 1.3.7 in [9]). Following Lemma 1.3, \mathcal{C} satisfies the split short five lemma. Since \mathcal{C} is monadic via the unit ball functor $U: \mathcal{C} \rightarrow \text{Set}$ (see [15], [20]) it is exact and thus semi-abelian. We are now going to give explicitly the operations witnessing this fact.

Following [18], in the theory of non-unital C^* -algebras we have the operations

$$\alpha_0(x, y) = \frac{1}{2}x - \frac{1}{2}y$$

and

$$\beta(x, y) = \bar{2}\bar{2}\left(\frac{1}{2}x + \frac{1}{4}y\right)$$

where

$$\bar{2} = 2(1 \vee |z|)^{-1}z$$

and

$$|z| = (z^* \cdot z)^{\frac{1}{2}}.$$

Then

$$\alpha_0(x, x) = \frac{1}{2}x - \frac{1}{2}x = 0$$

and

$$\beta(\alpha_0(x, y), y) = \beta\left(\frac{1}{2}x - \frac{1}{2}y, y\right) = \bar{2}\bar{2}\left(\frac{1}{4}x - \frac{1}{4}y + \frac{1}{4}y\right) = \bar{2}\bar{2}\left(\frac{1}{4}x\right) = \bar{2}\left(\frac{1}{2}x\right) = x.$$

1.5. Remark. The fact that commutative C^* -algebras form an exact Maltsev category was first observed in [8].

1.6. Example. The category $CompGrp$ of compact groups is monadic via the usual forgetful functor $U: CompGrp \rightarrow Set$. This immediately follows from the existence of free compact groups [11]. Since a group operation is present, $CompGrp$ is semi-abelian.

2. LOCALIZATIONS OF SEMI-ABELIAN VARIETIES

Let \mathcal{C} be a cocomplete category with a regular generator G and consider the functor $U: \mathcal{C}(G, -): \mathcal{C} \rightarrow Set$. There is a left adjoint F to U sending a set n to the n -th copower $n \cdot G$ of G . Let $T = UF$ be the induced monad and let $H: \mathcal{C} \rightarrow Alg(T)$ be the comparison functor. Since G is a regular generator, H is a full embedding. Theorem 1.1 tells us when $Alg(T)$ is semi-abelian. In terms of a generator G , the conditions in Theorem 1.1 can be expressed as follows:

- (1) there is exactly one arrow $0: G \rightarrow 0$ (where 0 is the initial object in \mathcal{C}).
- (2) there exist arrows $\alpha_i: G \rightarrow 2 \cdot G$, $i \in n$ (where $n \geq 1$ is a cardinal) such that the square

$$\begin{array}{ccc} G & \xrightarrow{\alpha_i} & 2 \cdot G \\ \downarrow & & \downarrow \nabla \\ 0 & \longrightarrow & G \end{array}$$

commutes (∇ is the codiagonal).

- (3) there is an arrow $\beta: G \rightarrow n \cdot G$ such that the diagram

$$\begin{array}{ccc} G & \xrightarrow{\beta} & n \cdot G \\ & \searrow i_1 & \swarrow (\alpha_0, \alpha_1, \dots, \alpha_i, \dots) \\ & 2 \cdot G & \end{array}$$

commutes (i_1 is the first injection into the coproduct).

We shall call a *semi-abelian generator* any regular generator G satisfying these three conditions. Of course, a free T -algebra $F(1)$ on 1 in a semi-abelian monadic category is a regular projective semi-abelian generator.

Now, let us recall that a full reflective subcategory \mathcal{C} of a category \mathcal{L} is a localization if the reflector $L: \mathcal{L} \rightarrow \mathcal{C}$ preserves finite limits.

2.1. Proposition. *A pointed category \mathcal{C} is a localization of a semi-abelian monadic category over Set if and only if \mathcal{C} is a cocomplete exact category with a semi-abelian generator.*

Proof. Due to the main result in [21] one already knows that localizations of monadic categories over Set are precisely cocomplete exact categories with a regular generator. More precisely, having such a category \mathcal{C} then the comparison functor $H: \mathcal{C} \rightarrow Alg(T)$ is a localization. This immediately yields the sufficiency, since the left adjoint L of H obviously preserves coproducts and the zero object, and then $L(F(1))$ is a semi-abelian generator in \mathcal{C} .

For the necessity, it suffices to observe that the functor H preserves the copowers of the semi-abelian generator G in \mathcal{C} . Consequently, $H(G)$ is a regular projective semi-abelian generator, and $Alg(T)$ is semi-abelian by Theorem 1.1. \square

2.2. Remark. There is an evident many-sorted version characterizing localizations of monadic categories over many-sorted sets as cocomplete exact categories having a semi-abelian generator in each sort.

2.3. Proposition. *A pointed category \mathcal{C} having copowers, pullbacks and a semi-abelian generator satisfies the split short five lemma.*

Proof. Under our assumption there is still a left adjoint F to $U = \mathcal{C}(G, -)$ and the comparison functor $H: \mathcal{C} \rightarrow Alg(T)$ is a full embedding. Since U preserves pullbacks, the forgetful functor $V: Alg(T) \rightarrow Set$ creates them and $V \circ H = U$, the functor H preserves pullbacks. Since $Alg(T)$ satisfies the split short five lemma, we conclude by Lemma 1.3 that \mathcal{C} satisfies it as well. \square

2.4. Remark. In order to give an example of a protomodular locally finitely presentable category \mathcal{C} with a zero object which does not have a semi-abelian generator, we will present it as an essentially algebraic theory Γ (see [1]). Let Γ contain a unique constant 0 , binary total operations α_0, γ and a binary partial operation $\beta(z, t)$ whose domain of definition $Def(\beta)$ is given by the equation $\gamma(z, t) = 0$. Let Γ contain the equations $\alpha_0(x, x) = 0$, $\gamma(\alpha_0(x, y), y) = 0$ and $\beta(\alpha_0(x, y), y) = x$. Then the argument used in Theorem 1.1 is still valid and $Alg(T)$ then satisfies the split short five lemma. On the other hand, β is not everywhere defined, which means that there is no reason for $Alg(T)$ to have a semi-abelian generator.

3. PROTOMODULAR QUASIVARIETIES

A category is quasimonadic over Set if it is a full regular epireflective subcategory of a monadic category over Set (i.e. a full reflective subcategory with the property that the unit of the adjunction is a regular epimorphism). Quasimonadic categories over Set are precisely cocomplete regular categories \mathcal{C} with a regular projective regular generator [10]. Again one uses $H: \mathcal{C} \rightarrow Alg(T)$ to present \mathcal{C} as a full regular epireflective subcategory of a monadic category.

3.1. Theorem. *A pointed category \mathcal{C} is a protomodular quasimonadic category over Set if and only if it is cocomplete, regular and has a regular projective semi-abelian generator.*

Proof. Every cocomplete regular category \mathcal{C} with a regular projective semi-abelian generator is a full regular epireflective subcategory of a semi-abelian monadic category $\mathit{Alg}(T)$ by Theorem 1.1. Hence it is pointed, quasimonadic and also satisfies the split short five lemma by Lemma 1.3.

Conversely, let \mathcal{C} be a regular epireflective subcategory of a semi-abelian monadic category $\mathit{Alg}(T)$, and let $F(1)$ be a free T -algebra on 1. Then the reflection of $F(1)$ to \mathcal{C} is a regular projective semi-abelian generator of \mathcal{C} . Moreover, \mathcal{C} is clearly cocomplete, regular and pointed. \square

In order to give a characterization of protomodular pointed quasivarieties, let us recall that an object G is *abstractly finite* if for any small set n there exists the n -th copower $S \cdot G$ of G and, moreover, any arrow $G \rightarrow S \cdot G$ factors through $S' \cdot G$ for some finite subset S' of S [13]. Any finitely presentable object is abstractly finite. Then from Corollary 4.4, Corollary 4.6 in [17] and Theorem 3.1 above the following results easily follow:

3.2. Corollary. *A pointed category \mathcal{C} is a regular epireflective subcategory of a protomodular finitary variety of universal algebras if and only if it is cocomplete, regular and has a regular projective abstractly finite semi-abelian generator.*

3.3. Corollary. *A pointed category \mathcal{C} is a protomodular quasivariety if and only if it is cocomplete, regular and has a finitely presentable regular projective semi-abelian generator.*

3.4. Example. The category Ab_{tf} of torsion-free abelian groups is an example of a pointed protomodular quasivariety. Indeed, Ab_{tf} is reflective in the category Ab of abelian groups, and it is closed in it under subobjects.

We conclude with the following

3.5. Theorem. *A pointed category \mathcal{C} is a localization of a protomodular quasimonadic category if and only if \mathcal{C} is a cocomplete regular category with a semi-abelian generator.*

Proof. Necessity is clear. Let then \mathcal{C} be a cocomplete regular category with a semi-abelian generator. Following the proof of Theorem 1.1 in [16], the category \mathcal{C} is a localization of its regular epireflective hull in $\mathit{Alg}(T)$. Then \mathcal{C} is a localization of a protomodular quasimonadic category over Set . \square

All the results in this section have obvious many-sorted versions.

REFERENCES

- [1] J. Adámek and J. Rosický, *Locally Presentable and Accessible Categories*, Lond. Math. Soc. Lect. Notes Ser. 189, Cambridge University Press, 1994.
- [2] M. Barr, *Exact categories*, Springer Lecture Notes Math., 236, 1971, 1-120.
- [3] F. Borceux, *A survey on semi-abelian categories*, to appear in the *Fields Institute Communications Series*.
- [4] D. Bourn, *Normalization equivalence, kernel equivalence and affine categories*, Springer Lecture Notes Math. 1488, 1991, 43-62.

- [5] D. Bourn, *Normal subobjects and abelian objects in protomodular categories*, Journal of Algebra, 228, 2000, 143-164.
- [6] D. Bourn, *3×3 Lemma and Protomodularity*, Journal of Algebra, 236, 2001, 778-795.
- [7] D. Bourn and G. Janelidze, *Characterization of protomodular varieties of universal algebras*, Th. Appl. Categories, 11,6 2003, 143-147.
- [8] A. Carboni, G.M. Kelly and M.C. Pedicchio, *Some remarks on Maltsev and Goursat categories* Appl. Categorical Structures, 1, 1993, 385-421.
- [9] J. Dixmier, *Les C^* -algèbres et leurs représentations*, Gauthier-Villars Editeurs, 1969.
- [10] W. Felscher, *Kennzeichnung von primitiven und quasi-primitiven Kategorien von Algebren*, Arch. Math. 39, 1968, 390-397.
- [11] K. Hoffmann *An essay on free compact groups*, Categorical aspects of topology and analysis, Proc. int. Conf. Ottawa 1980, Lect. Notes Math. 915, 1982, 171-197.
- [12] G. Janelidze, L. Marki and W. Tholen, *Semi-Abelian categories*, J. Pure Appl. Algebra 168, 2002, 367-386.
- [13] F.W. Lawvere, *Functorial semantics of algebraic theories*, Proc. Nat. Acad. Sc. USA 50, 1963, 869-872.
- [14] E.G. Manes, *Algebraic Theories*, Springer-Verlag, 1976.
- [15] J.W. Negrepointis, *Duality in analysis from the point of view of triples*, Journal of Algebra, 19, 1971, 228-253.
- [16] M.C. Pedicchio and J. Rosický, *Localizations of varieties and quasivarieties*, J. Pure Appl. Algebra 148, 2000, 275-284.
- [17] M.C. Pedicchio and E.M. Vitale, *On the abstract characterization of quasi-varieties*, Alg. Universalis 43, 2000, 269-278.
- [18] J. Pelletier and J.Rosický, *On the equational theory of C^* -algebras*, Alg. Universalis 30, 1993, 275-284.
- [19] A. Ursini, *On subtractive varieties*, Alg. Universalis 31, 1994, 204-222.
- [20] D. Van Osdol, *C^* -algebras and cohomology*, Proc. Conf. Categorical Topology, Toledo 1983, Heldermann Verlag, 1984, 582-587.
- [21] E.M. Vitale, *Localizations of algebraic categories*, J. Pure Appl. Algebra 108, 1996, 315-320.

UNIVERSITÉ DU LITTORAL CÔTE D'OPALE, LABORATOIRE DE MATHÉMATIQUES PURES ET APPLIQUÉES, BT. H. POINCARÉ, 50 RUE F. BUISSON BP 699, 62228 CALAIS, FRANCE.

MASARYK UNIVERSITY, DEPARTMENT OF MATHEMATICS, JANÁCKOVO NÁM. 2A, 662 95 BRNO, CZECH REPUBLIC.

E-mail address: gran@lmpa.univ-littoral.fr and rosicky@math.muni.cz