

We will continue online on Thursday, **May 13th, at 13.00 CEST** on [ZOOM](#) platform (for information how to access seminar and next programme visit

[this page](#)

) by the talk:

Nathanael Arkor

Higher-order algebraic theories and relative monads

Abstract:

There have traditionally been two ways to reason about universal algebraic structure categorically: via algebraic theories, and via monads. It is well known that the two are tightly related: in particular, there is a correspondence between algebraic theories and a class of monads on the category of sets.

Motivated by the study of simple type theories, Fiore and Mahmoud introduced second-order algebraic theories, which extend classical (first-order) algebraic theories by variable-binding operators, such as the existential quantifier $\exists x$ of first-order logic; the differential operators d/dx analysis; and the λ -abstraction operator of the untyped λ -calculus. Fiore and Mahmoud established a correspondence between second-order algebraic theories and a second-order equational logic, but did not pursue a general understanding of the categorical structure of second-order algebraic theories. In particular, the possibility of a monad–theory correspondence for second-order algebraic theories was left as an open question.

In this talk, I will present a generalisation of algebraic theories to higher-order structure, in particular subsuming the second-order algebraic theories of Fiore and Mahmoud, and describe a universal property of the category of n th-order algebraic theories. The central result is a correspondence between $(n + 1)$ th-order algebraic theories and a class of relative monads on the category of n th-order algebraic theories, which extends to a monad correspondence subsuming that of the classical setting. Finally, I will discuss how the perspective lent by higher-order algebraic theories sheds new light on the classical monad–theory correspondence.

This is a report on joint work with Dylan McDermott.